

RESEARCH

Open Access



Integral representations of equally positive integer-indexed harmonic sums at infinity

Lin Jiu

*Correspondence:
 ljiu@risc.uni-linz.ac.at
 Research Institute for Symbolic
 Computation, Johannes Kepler
 University, 4040 Linz, Austria

Abstract

We identify a partition-theoretic generalization of Riemann zeta function and the equally positive integer-indexed harmonic sums at infinity, to obtain the generating function and the integral representations of the latter. The special cases coincide with zeta values at positive integer arguments.

Keywords: Harmonic sum, Integral representation, Zeta value

1 Background

The harmonic sum of indices $a_1, \dots, a_k \in \mathbb{R} \setminus \{0\}$ is defined as (see [1, eq. 4, pp. 1])

$$S_{a_1, \dots, a_k}(N) = \sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} \frac{\text{sign}(a_1)^{n_1}}{n_1^{|a_1|}} \times \dots \times \frac{\text{sign}(a_k)^{n_k}}{n_k^{|a_k|}},$$

which is naturally connected to the Riemann zeta function, by noting that $N = \infty, k = 1$ and $a_1 > 0$ gives $S_{a_1}(\infty) = \zeta(a_1)$. A variety of the study can be found in the literature. For instance, Hoffman [4] established the connection between harmonic sums and multiple zeta values. We especially focus on the *equally positively indexed harmonic sums*, given by the case $a_1 = \dots = a_k = a > 0$

$$S_{a_k}(N) := S_{\underbrace{a, \dots, a}_k}(N) = \sum_{N \geq n_1 \geq \dots \geq n_k \geq 1} \frac{1}{(n_1 \dots n_k)^a}, \tag{1.1}$$

and also the *equally positive integer-indexed harmonic sums* (EPIIHS), namely $a = m \in \mathbb{Z}_{>0}$. If $N = \infty$, we additionally assume $m \in \mathbb{Z}_{>1}$ for convergence.

Recently, Schneider [6] studied the generalized q -Pochhammer symbol and obtained [6, pp. 3]

$$\prod_{n \in X} \frac{1}{1 - f(n)q^n} = \sum_{\lambda \in \mathcal{P}_X} q^{|\lambda|} \prod_{\lambda_i \in \lambda} f(\lambda_i), \tag{1.2}$$

where

- $X \subseteq \mathbb{Z}_{>0}$ and $f : \mathbb{Z}_{>0} \rightarrow \mathbb{C}$ such that if $n \notin X$ then $f(n) = 0$;
- \mathcal{P}_X is the set of partitions into elements of X ;
- $\lambda \vdash n$ means λ is a partition of n , the size $|\lambda|$ is the sum of the parts of λ , i.e., the number n being partitioned, and $\lambda_i \in \lambda$ means $\lambda_i \in \mathbb{Z}_{>0}$ is a part of partition λ .

© The Author(s) 2017. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Further define $l(\lambda) := k$, $n_\lambda := \lambda_1 \cdots \lambda_k$ and denote $\mathcal{P} := \mathcal{P}_{\mathbb{Z}_{>0}}$. Noting $\lambda_1 \geq \cdots \geq \lambda_k \geq 1$, a *partition-theoretic generalization of Riemann zeta function* [6, eq. 11, pp. 4] is defined and identified as

$$\zeta_{\mathcal{P}}(\{a\}^k) := \sum_{l(\lambda)=k} \frac{1}{n_\lambda^a} = \sum_{\lambda_1 \geq \cdots \geq \lambda_k \geq 1} \frac{1}{\lambda_1^a \cdots \lambda_k^a} = S_{a_k}(\infty), \tag{1.3}$$

which leads to the generating function and the integral representation of $S_{m_k}(\infty)$, presented in the next section.

2 Main results

We first apply (1.2) to the case $X = \{1, 2, \dots, N\}$ and $f(n) := \frac{t^a}{n^a}$, obtaining

$$\prod_{n=1}^N \frac{1}{1 - \frac{t^a}{n^a} q^n} = \sum_{\lambda \in \mathcal{P}_X} q^{|\lambda|} \prod_{\lambda_i \in \lambda} \frac{t^a}{\lambda_i^a} = \sum_{\lambda \in \mathcal{P}_X} q^{|\lambda|} \frac{t^{l(\lambda)a}}{n_\lambda^a},$$

which, by further letting $q \rightarrow 1$, yields the following generating function.

Theorem 1 *The generating function of $S_{a_k}(N)$ is given by*

$$\sum_{k=0}^{\infty} S_{a_k}(N) t^{ak} = \prod_{n=1}^N \frac{n^a}{n^a - t^a}. \tag{2.1}$$

Remark 2 The special case for $a = 1$ is [2, eq. 9, pp. 1272]

$$\sum_{k=0}^{\infty} t^k S_{1_k}(N) = \frac{N!}{(1-t) \cdots (N-t)} = N \cdot B(N, 1-t), \tag{2.2}$$

involving the beta function B , defined by

$$B(x, y) := \int_0^1 z^{x-1} (1-z)^{y-1} dz = \frac{\Gamma(x) \Gamma(y)}{\Gamma(x+y)}, \tag{2.3}$$

where the integral representation holds for $\text{Re}(x), \text{Re}(y) > 0$.

Corollary 3 *For $m \in \mathbb{Z}_{>1}$, denote $\xi_m := \exp\left(\frac{2\pi i}{m}\right)$ with $i^2 = -1$. Then,*

$$\sum_{k=0}^{\infty} S_{m_k}(\infty) t^{mk} = \prod_{j=0}^{m-1} \Gamma\left(1 - \xi_m^j t\right). \tag{2.4}$$

Proof From (2.1) and (2.2), we have

$$\sum_{k=0}^{\infty} S_{m_k}(N) t^{mk} = \prod_{n=1}^N \frac{n^m}{(n - \xi_m^0 t) \cdots (n - \xi_m^{m-1} t)} = \prod_{j=0}^{m-1} N \cdot B\left(N, 1 - \xi_m^j t\right).$$

Then, apply the limit (see [7, pp. 254, ex. 5]) $\Gamma(z) = \lim_{N \rightarrow \infty} N^z B(N, z)$ to $z_j = 1 - \xi_m^j t$, $j = 0, \dots, m-1$, by noting $\xi_m^0 + \cdots + \xi_m^{m-1} = 0$, to complete the proof. \square

Remark 4 An alternative proof can be given by letting $N = \infty$ in (2.1) and applying [3, Thm. 1.1, pp. 547].

Remark 5 For general $a > 0$, we failed to obtain a closed form of $\prod_{n=1}^{\infty} \frac{n^a}{n^a - t^a}$.

Example 6 When $m = 2$, we apply (2.4) to get

$$B(1+t, 1-t) = \Gamma(1+t)\Gamma(1-t) = \sum_{k=0}^{\infty} S_{2k}(\infty) t^{2k}.$$

From the integral representation (2.3), we obtain (also see Remark 7)

$$B(1+t, 1-t) = \int_0^1 z^t (1-z)^{-t} dz = \sum_{k=0}^{\infty} \frac{t^k}{k!} \int_0^1 \log^k \left(\frac{z}{1-z} \right) dz. \tag{2.5}$$

Then it follows, by comparing coefficients of t ,

$$S_{2k}(\infty) = \frac{1}{(2k)!} \int_0^1 \log^{2k} \left(\frac{z}{1-z} \right) dz.$$

In particular, $k = 1$ yields

$$\frac{\pi^2}{6} = \zeta(2) = S_2(\infty) = \frac{1}{2} \int_0^1 \log^2 \left(\frac{z}{1-z} \right) dz.$$

Remark 7 We may interchange the integral and the sum of the series in (2.5), by restricting t to a closed compact set, e.g., $[-\frac{1}{2}, \frac{1}{2}]$, satisfying $\text{Re}(1-t), \text{Re}(1+t) > 0$ as that in (2.3), in order to guarantee uniform convergence of the integral representation. (Similar discussion is omitted for the multiple beta function, defined next.)

Definition 8 The *multiple beta function* [5, Ch. 49] is defined as

$$B(\alpha_1, \dots, \alpha_m) := \frac{\Gamma(\alpha_1) \cdots \Gamma(\alpha_m)}{\Gamma(\alpha_1 + \cdots + \alpha_m)} = \int_{\Omega_m} \prod_{i=1}^m x_i^{\alpha_i - 1} d\mathbf{x}, \tag{2.6}$$

where $\Omega_m = \{(x_1, \dots, x_m) \in \mathbb{R}_{>0}^m : x_1 + \cdots + x_{m-1} < 1, x_1 + \cdots + x_m = 1\}$ and the integral representation requires $\text{Re}(\alpha_1), \dots, \text{Re}(\alpha_m) > 0$.

Following the same idea as that in Example 6, we first have, from (2.4),

$$B(1 - \xi_m^0 t, \dots, 1 - \xi_m^{m-1} t) = \frac{1}{(m-1)!} \sum_{k=0}^{\infty} S_{m_k}(\infty) t^{mk}.$$

Then, apply the integral representation (2.6), expand the integrand as a power series in t , and compare coefficients of t , to obtain the following integral representation.

Theorem 9 For all $m, k \in \mathbb{Z}_{>0}$ with $m \geq 2$,

$$S_{m_k}(\infty) = \frac{(-1)^{mk} (m-1)!}{(mk)!} \int_{\Omega_m} \log^{mk} \left(\prod_{j=0}^{m-1} x_{j+1}^{\xi_m^j} \right) d\mathbf{x}.$$

Corollary 10 In particular, the case $k = 1$ implies for integer $m \in \mathbb{Z}_{>1}$ that

$$\zeta(m) = \frac{(-1)^m}{m} \int_{\Omega_m} \log^m \left(\prod_{j=0}^{m-1} x_{j+1}^{\xi_m^j} \right) d\mathbf{x},$$

or alternatively

$$\zeta(m) = \frac{(-1)^m}{m} \int_0^1 \int_0^{1-x_1} \dots \int_0^{1-x_1-\dots-x_{m-2}} \log^m \left(x_1^{\xi_m^0} \dots x_{m-1}^{\xi_m^{m-2}} (1-x_1-\dots-x_{m-1})^{\xi_m^{m-1}} \right) \times dx_{m-1} \dots dx_1.$$

Acknowledgements

The author would like to thank Dr. Jakob Ablinger for his help on harmonic sums; Prof. Johannes Blümlein for his handwritten notes on the proof of (2.2); and especially his mentors, Prof. Peter Paule and Prof. Carsten Schneider, for their valuable suggestions. This work is supported by the Austrian Science Fund (FWF) Grant SFB F50 (F5006-N15 and F5009-N15).

Received: 14 November 2016 Accepted: 25 January 2017

Published online: 10 May 2017

References

1. Blümlein, J., Kurth, S.: Harmonic sums and Mellin transforms up to two-loop order. *Phys. Rev. D* **60**, 014018 (1999)
2. Blümlein, J., Klein, S., Schneider, C., Stan, F.: A symbolic summation approach to Feynman integral calculus. *J. Symb. Comput.* **47**, 1267–1289 (2012)
3. Chamberland, M., Straub, A.: On gamma quotients and infinite products. *Adv. Appl. Math.* **51**, 546–562 (2013)
4. Hoffman, M.: Multiple harmonic series. *Pac. J. Math.* **152**, 275–290 (1992)
5. Kotz, S., Balakrishnan, N., Johnson, N.L.: *Continuous Multivariate Distributions Vol 1: Models and Applications*. Wiley, New York (2010)
6. Schneider, R.: Partition zeta functions. *Res. Number Theory* **2**, 9 (2016)
7. Whittaker, E.T., Watson, G.N.: *A Course of Modern Analysis*. Cambridge Univ. Press, New York (1965)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
